MONOCLONAL ANTIBODIES (MABs) – ANTI-IL, ANTI-IgE, ANTI-TSLP PRIOR AUTHORIZATION FORM





(form effective 1/8/2024)

Fax to PerformRx[™] at **1-888-981-5202**, or to speak to a representative call **1-866-610-2774**.

PRIOR AUTHORIZATION REQUEST INFORMATION							
	enewal request	Total # of pages:					
Name of office contact:			Contact's phone number:		LTC fa	LTC facility contact/phone:	
PATIENT INFORMATION							
Patient name: Patient ID #:						DOB:	
						565.	
Street address: Apt #: City/state/zip: Phone:							
PRESCRIBER INFO	ORMATION						
Prescriber name:				NDI		State liegenes #	
Specialty:				NPI:		State license #:	
Street address:							
Suite #:	City/state/zip:						
Phone:				Fax:			
CLINICAL INFORM	1ATION						
Medication requested:						Strength:	
Preferred Medications:			No	Non-Preferred Medications:		Dosage form (pen, vial, etc):	
🗆 Fasenra Pen		Tezspire Pen		Cinqair Vial			
Fasenra Syringe		Xolair Syringe		Nucala 100 mg/ml Syringe			
				Nucala 100 mg/ml Vial			
Nucala 40 mg/0.4 ml Syringe Tezspire Syringe							
Dose and directions:				iantity:		Duration: months	
Diagnosis: Dx code <u>(required)</u> :						Weight: Ibs/kg	
Has the beneficiary used the requested medication in the past 90 days? Submit documentation.						\Box Yes – date of last dose:	
						□ No	
Is the requested medication being prescribed by or in consultation with a specialist?						□ Yes Submit documentation of □ No consultation, if applicable.	
PHARMACY INFORMATION (Prescriber to identify the pharmacy that is to dispense the medication):							
Deliver to: Patient's Home Physician's Office Patient's Preferred Pharmacy Name:							
NPI#:							
Pharmacy Phone #: Pharmacy Fax #:							
□ I acknowledge that the patient agrees with the pharmacy chosen for delivery of this medication.							
INITIAL REQUESTS							
Complete all sections that apply to the beneficiary and this request. Check all that apply and <u>submit documentation</u> for each item.							
For a non-preferred drug in this class: Does the beneficiary have a history of trial and failure of or contraindication or an intolerance to the preferred agents in this class that are approved or medically accepted for treatment of the beneficiary's condition? Refer to https://papdl.com/preferred-drug-list for a list of preferred and non-preferred agents in this class.					e to the	□ Yes Submit documentation. □ No	
1. For treatment of ASTHMA: I Has an asthma severity that is consistent with the FDA-approved indication for the requested medication despite use of maximal therapeutic doses of or has contraindication or an intolerance to the following (check all that apply): I inhaled glucocorticoid I long-acting beta-agonist (LABA) I eukotriene modifier For an anti-IgE MAB (e.g., XOLAIR): I Has moderate-to-severe persistent asthma induced by an unavoidable per allergen (pollen, mold, dust mites, etc.) I biggious confirmed by positive skin test or radioallergosorbent test (RAST I eukotriene modifier							
🗆 other (e.g., tiotro	pium, theophylline): maximal standard asthr	sthma controller medications in add	lition	Has asthma of an eosinopl	IL MAB (e.g., CINQAIR, FASENRA, NUCALA): na of an eosinophilic phenotype – Absolute blood eosinophil count: /mL Date obtained: e asthma		
				□ For an anti-TSLP (e.g., TEZS □ Has severe asthma			



INITIAL REQUESTS (continued)

2. For treatment of CHRONIC SPONTANEOUS (IDIOPATHIC) URTICARIA:

- \Box Has a history of urticaria for a period of ≥ 6 weeks
- □ Requires use of systemic steroids to control urticarial symptoms
- Tried and failed the maximally tolerated dose of an H1 antihistamine (e.g., cetirizine/levocetirizine, fexofenadine, loratadine/desloratadine) taken for at least two weeks or has a contraindication or an intolerance to H1 antihistamines
- 3. For treatment of EGPA:
 - \Box Has a history of asthma
 - □ Has an absolute blood eosinophil count ≥1000/microliter
 - \Box Has a blood eosinophil level >10% of leukocytes
 - □ Has evidence of the following (check all that apply):
 - □ histopathological evidence of:
 - eosinophilic vasculitis
 - perivascular eosinophilic infiltration
 - $\hfill\square$ eosinophil-rich granulomatous inflammation
 - neuropathy (nerve deficit or conduction abnormality)
 - pulmonary infiltrates, non-fixed
 - □ sino-nasal abnormality
 - □ cardiomyopathy
 - □ glomerulonephritis
 - □ alveolar hemorrhage
 - □ palpable purpura
 - □ positive test for ANCA
 - Requires systemic glucocorticoids to maintain remission
 - Has a contraindication or an intolerance to systemic glucocorticoids
 - □ Has severe EGPA as defined by national treatment guidelines
 - □ Tried and failed or has a contraindication or an intolerance to rituximab or cyclophosphamide

4. For treatment of HYPEREOSINOPHILIC SYNDROME (HES):

- □ Has documented FIP1L1-PDGFRA-negative HES
- □ Has organ damage or dysfunction
- □ Has a blood eosinophil count ≥1000/microliter
- □ Requires or has required systemic glucocorticoids to maintain remission
- □ Has a contraindication or an intolerance to systemic glucocorticoids

5. For treatment of NASAL POLYPS:

- Has a history of trial and failure of or contraindication or intolerance to nasal corticosteroids
- □ For an anti-IgE MAB (e.g., XOLAIR):
 - Has a pretreatment serum total IgE measurement of: _____
- 6. For treatment of ALL OTHER DIAGNOSES:
 - List other treatments tried (including start/stop dates, dose, outcomes):

RENEWAL REQUESTS

1. For treatment of ASTHMA:

- Experienced measurable evidence of improvement in the severity of the asthma condition
- □ Will continue to use optimally titrated doses of or has a contraindication or an intolerance to the following (check all that apply):
 - □ inhaled glucocorticoid
 - □ leukotriene modifier
 - □ long-acting beta-agonist (LABA)
 - □ other (e.g., tiotropium, theophylline):

2. For treatment of CHRONIC SPONTANEOUS (IDIOPATHIC) URTICARIA:

- □ Experienced an improvement in symptoms
- □ Document rationale for continued use: ____

3. For treatment of EGPA:

□ Experienced measurable evidence of improvement in disease activity □ Reduction in use of systemic glucocorticoids for the treatment of EGPA

4. For treatment of HYPEREOSINOPHILIC SYNDROME (HES):

- Experienced measurable improvement in disease activity
 - Reduction in use of systemic glucocorticoids for the treatment of HES

PLEASE FAX COMPLETED FORM WITH REQUIRED CLINICAL DOCUMENTATION

Prescriber signature:

Confidentiality Notice: The documents accompanying this telecopy may contain confidential information belonging to the sender. The information is intended only for the use of the individual named above. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution or taking of any telecopy is strictly prohibited.

Date: